How Your Medication Error Happened, and How You Plan To Keep It From Happening Again

| Date of medication err | or: |
|------------------------|-----|
| UCI #: | |
| SANDIS Incident #: | |

| What went wrong with the individual's medications (meds)? (Please check all that apply.) | | | | |
|-------------------------------------------------------------------------------------------------------------------|--|--|--|--|
| Missed dose Wrong med(s) Wrong dose Wrong Route Wrong person Wrong time | | | | |
| Who was tasked to give the meds? (Please check one.) Nurse Other Staff Family Self | | | | |
| Giving/taking meds happens in steps: | | | | |
| | | | | |
| Step 1: Step 2: Step 3: | | | | |
| Set-up Give/Take Meds Record/Check | | | | |
| | | | | |
| What step(s) went wrong and how did they go wrong? (Please check all that apply.) | | | | |
| Step 1: Set-up/Prepare Meds | | | | |
| | | | | |
| Did not start on time. | | | | |
| Did not check med log before getting meds from storage. | | | | |
| Did not compare labels on meds with med log when getting meds from storage. | | | | |
| Meds in log not the same as current physician orders. Meds in log not the same as labels on bottles, packaging. | | | | |
| Took wrong meds from storage: misread label or med log. | | | | |
| Did not get all meds: left some meds in storage. | | | | |
| Not able to find some meds: meds not in usual place. | | | | |
| Not able to find some meds: meds not in usual place. Not able to find some meds: meds not ordered or re-ordered. | | | | |
| Not able to find some meds: pharmacy failed to fill some ordered meds. | | | | |
| Physician orders not copied correctly by the pharmacy or in med log. | | | | |
| Individual was away from home and did not have meds along. | | | | |
| Step 2: Give/Take Meds | | | | |
| Did not place all meds in the cup when pouring. | | | | |
| Did not give all of the meds in the cup or bubble pack. | | | | |
| Did not give meds to the correct individual. | | | | |
| Was interrupted while giving meds; made an error when resumed. | | | | |
| Dropped or lost a pill when giving it, or Individual dropped or lost a pill when taking it. | | | | |
| Individual was not able to follow the medication routine as planned. | | | | |
| Step 3: Record/Check Meds Afterward | | | | |
| Staff who gave meds did not log meds that an individual took. | | | | |
| Staff who gave meds did not log time when an individual took meds. | | | | |
| Staff who gave meds did not check meds, packs, and log after. | | | | |
| No one else checked meds, packs after. | | | | |
| No one else checked the med log after. | | | | |
| Almost done! Please turn page over. | | | | |

| Date of medication error: | |
|---------------------------|--|
| UCI #: | |
| SANDIS Incident #: | |

| Besides training, what will you do to stop this from happening again? | | | | |
|-----------------------------------------------------------------------------------------------|---------------------------------------|----------------------------------------------------|--|--|
| (Please check all that apply.) | | | | |
| Step 1: Set-up/Prepare Meds | | | | |
| Have a way to alert staff and individual when it is time for meds. | | | | |
| Use a med log with simple, clear design that is easy to read. | | | | |
| Assign someone to order meds and check that pharmacy filled order. | | | | |
| Assign someone to check med log and physician orders, and update the log. | | | | |
| Assign someone to check med log and labels on bottles and packaging, and update the log. | | | | |
| Assign back-up staff to order, prepare, and give meds if regular staff are absent. | | | | |
| Organize storage in a simple, clear way. | | | | |
| Store meds in a space that is well lit, where meds are easy to see. | | | | |
| Ask pharmacy for their most curre | ent med log. | | | |
| Assign someone to check that Individuals have meds when away. | | | | |
| Step 2: Give/Take Meds | | | | |
| If pouring meds, have staff check | meds against med log as they place | them in cup. | | |
| Have staff indicate each med that they give in the med log, and when they gave it. | | | | |
| Have staff perform a gloved finger | r swipe of bubble packs after giving | meds to an individual. | | |
| If someone is giving meds, do not allow him/her to be interrupted. | | | | |
| Designate a place for preparing ar | nd giving meds. | | | |
| Give meds to one individual at a ti | ime, away from other people and ac | tivities. | | |
| Give pills across a table, to decrease risk of dropped meds. | | | | |
| Plan routines with Individuals: Ask | them how they want the routine to | go. | | |
| Change routines that are not work | king (e.g. ask if doctor can change p | rescribed times). | | |
| Step 3: Record/Check Meds Afterwar | rd | | | |
| Use a med log with simple, clear of | lesign and that is easy to read, upda | te. | | |
| Assign someone to check med log and physician orders, and update the log. | | | | |
| Assign someone to check the med log against labels on bottles, packaging, and update the log. | | | | |
| Use bubble packs. | | | | |
| Check bubble packs after giving meds by a gloved finger swipes. | | | | |
| Have a second person check bubble packs by a second gloved finger swipe. | | | | |
| Have a second person check med log. | | | | |
| Have a plan for using a med log and updating it regularly. | | | | |
| Date when you plan to have your changes in place: | | | | |
| | | | | |
| Person completing this document: | | | | |
| Vendor Name | Vendor Number | | | |
| | | Developed by Mission Analytics Group, Inc. for the | | |
| Comments: | | California Department of Developmental Services | | |
| | You are done! Thank you 😊 | www.missionanalyticsgroup.com | | |